

Evaluation of pH Levels in Some Common Carbohydrate Food Items Consumed by Communities in the Central Senatorial District of Cross River State, South-South of Nigeria

E.N. Tawo¹, A.E. Abara¹, S.P. Malu² and N.O. Alobi¹

¹Department of Chemical Sciences, Cross River University of Technology, Calabar, Nigeria

²Department of Chemistry, Benue State University, Makurdi, Benue State, Nigeria

Abstract: This work evaluates and provides data on the seasonal pH of common carbohydrate food sources in major communities of central senatorial districts of Cross River State. The result shows that most of the locally consumed food items within these zones are mostly acidic within a mean value of pH of 4.6 and conductivity of 48.6 Siemens for the wet season and pH of 4.8 and conductivity of 27.4 for the dry season which causes a lot of ailments. A balance diet or a combination of food substances to keep an optimal body pH balance is recommended so as to discourage the growth of microforms and organisms which give rise to these problems. It is recommended also that other non-carbohydrate food items such as proteins and vegetables be increased in the menu, or the starchy content of food substances can be reduced during the manufacturing process by manufacturing companies. Government can legislate and monitor these companies to comply with such standards as to improve the health conditions of the rural communities and other consumers.

Key words: Acidity, alkalinity, pH, conductivity, carbohydrate foods

INTRODUCTION

Common food items like Garri, Yam, Cocoyam, Cassava, Plantain are rich in carbohydrates; meat, fish, beans are fairly rich in proteins; the cereals and legumes like rice ground nuts, peanuts, cashew, contain appropriate fats and oils; vegetables and fruits contain trace metals and vitamins. A balanced diet contains all these in their correct ratios. A combination of these food items in their correct amounts guarantees a fair optimal pH balance of the body. Which aids the elimination of such maladies as pain, arthritis, cramps, diabetes mellitus, respiratory problems, anti social behaviour and suicidal tendencies among others (Susan, 2000).

Carbohydrate foods such as cassava, yam, cocoyam, plantain, bananas, maize in their raw form, semi-processed and processed forms like garri, cassava flour, yam flour (amala), maize flour, are a common feature of the diet of most Nigerian communities. These foods are acidic. There is a general tendency among living organisms especially humans, towards the consumption of acidic food items. (Susan, 2000; Young and Young, 2002). However the human race, in its quest for longevity, struggles, just as the cells do, to maintain a safe pH balance until death (Halprin and Goldstein, 1999).

Some materials have been in circulation dealing with healthy living through a conscious pH balance. Chemically, this is a struggle for adequate pH control of the aging process which is a natural process of acid generation and acid waste breakdown and disposal (Halprin and Goldstein, 1999; Young and Young, 2002).

The feeding habits of people within a community dictate to a large extent their body pH balance and the food they eat is the source of the ingredients that facilitate the metabolic processes that lead to the generation and breakdown of acidic/basic by products (Hall, 1976; Remer and Manz, 1994)

The combination therefore, of the various food substances to check this imbalance depends on the availability of alternative food and the purchasing power of the people, including of course their traditional beliefs. (Oyenuga, 1968; Hall, 1976)

Cross River State like most States in Nigeria is "an assemblage" of peasant communities that rely mostly on carbohydrate/starchy foods for their sustenance. The soil that supports the growth of these food substances contributes the trace metals which are a necessary accompaniment of their menu (Eneobong, 2001). These metals cause the hydrolysis of foods to form acidic or basic end-products which accumulate in body cells during various metabolic processes (Hall, 1976; Harris, 1975; Susan, 2000).

This research therefore is aimed at building up data on pH levels in some common food items available within our local communities (Anon, 1962; Susan, 2000), as a base of information to aid a balanced diet in order to achieve an optimal body pH balance, which is said to be a remedy for many body ailment, common to the people in our local communities (Green and Kleeman, 1991; Young and Young, 2002).

The developed nations of the world, like the United States of America have enough of this type of data

management, for their food items and balanced diet. However, when they recommend recipes to third world countries like ours, we are at a loss as to what their local equivalents are. Here lies the yearning for such data and justification for this work, which cannot be overemphasized (Young and Young, 2002).

MATERIALS AND METHODS

The raw food items, the semi-processed and processed food items were collected from the Central Senatorial District of Cross River State which comprises Abi, Yakurr, Obubra, Ikom, Etung and Boki Local Government Areas. The collection centres were grouped into three Stations, namely Akparabong in Ikom Local Government Area as Station I, Boki in Boki Local Government Area as Station II and Ugep in Yakurr Local Government Area as Station III.

The Raw Food items include Cassava, Yam, Cocoyam, Sweet/Bitter Yam, Rice, Banana, Plantain, Maize/Corn, Sesame, Sugar Cane, Mango, Grape Fruit, Lemon, Cucumber, Paw paw, Coconut, Avocado. While the Semi-Processed/Processed food items include Garri, Fufu, Bread, Yam flour (Amala), Cassava flour, Wheat flour, Plantain flour and Spaghetti/Macarroni, Semovita/Semolina.

All samples were collected at six points in a rectangular grid method in each of these stations to ensure good representation (Osuala, 1980).

Samples collected from each station were thoroughly mixed to form a composite sample. About 12 raw samples were collected to include cucumber, plantain, rice, maize, banana, pineapple, yam, tomato, cassava, grape, orange, mango and potato.

Treatment of samples: For raw yam, cassava, potato, rice, banana, plantain and cocoyam, peeling was done for each using a properly cleaned kitchen knife. About 10 g of each sample was cut to small pieces and placed into a 1 litre-cup Kenwood Model no KW10 blender and 30 ml of water added. The samples were then pulverized for 5 min. Subsequent samples were treated the same way after thorough cleaning of the blender.

For oranges, grapes and other citrus fruits, the liquid-juice was squeezed into beakers and their concentrated juice covered and kept for analysis.

For the semi-processed and processed food items like cassava flour, Akpu and Garri, about 10 gm of each sample was weighed using a Mettler Top-loading balance and placed into separate beakers. 100 ml of distilled water was added to each of them and stirred and kept for sample analysis. Care was taken to ensure that the collected samples were not wounded to avoid any fast enzyme activity. For samples like cassava flour and Akpu, the cassava was grated and fermented for two days after which they were washed with water and later sieved to collect the wet Akpu samples. The other

fermented portion was washed and then dried using sunlight and ground to form the cassava flour (Akinrele, 1976; Eka, 1986; Achinewhu, 1998)

Preparation of garri: The cassava tubers was peeled and properly washed. It was then grated and tied overnight in sacco bag for some of the water to drain off. The pressed dry portion was loosened and sifted to remove the unwanted fibers. It was now ready for frying with locally-made ceramic frying pan. (Akinrele, 1976; Eka, 1986).

Laboratory analysis of samples: A digital pH meter, pH indicator papers and a conductivity meter Model No 4510 were used to determine acidity. Due to seasonal variations, determinations were carried out in two phase. The first phase covered the dry season (November to April) while the second phase covered the rain season (May to October). The indicator papers and conductivity measurement formed the preliminary test for determining acidity, digital pH meter gave the precise results (Krist and Rump, 1988).

About 10 gm of each sample was weighed into three 250 ml beakers; 100 ml of distilled water was added. The sample was stirred for 5 min with a clean glass rod and allowed to equilibrate. The sample was then covered with a watch glass to avoid exposure to the atmosphere and kept overnight.

The supernatant liquid in the first beaker was tested using pH indicator paper to give the approximate pH. The solution in the second beaker was tested for conductivity by placing the dip cell of the meter into it at the ambient temperature. In the third beaker the pH was determined electronically using a pH meter of +0.1 percent sensitivity Model No WTW 422. The instrument was carefully calibrated using Buffers of pH 4, pH 7 and pH 9 (Krist and Rump, 1988). The use of filter papers to blot the electrode after each calibration was meticulously followed to reduce memory effect. The pH of the samples was thus determined and the results are shown in Tables 1 and 2 for the rainy season and dry season period.

RESULTS AND DISCUSSION

The seasonal mean value of the pH of common carbohydrate food items is shown in Tables 1 and 2. the result shows that the mean value of pH of 4.6 and conductivity of 48.6 Siemens was obtained for rainy season while that of dry season was pH of 4.8 and conductivity of 27.4 Siemens. It can be inferred from these results that carbohydrate food substances generate acidic waste products (Susan, 2000).

By design the human body is alkaline but the intake of carbohydrates either in their raw state or processed form renders them ultimately acidic. Available evidence shows that this tendency on increasing body acidity by

Table 1: Rainy Season May-October Mean pH and Conductivity in Siemens ± standard deviation

Sample	Mean P ^H	±	STD	Conductivity	±	STD
Cucumber	4.31	±	0.01	49.8	±	0.3
Plantain	4.15	±	0.01	52.9	±	0.2
Rice (white)	6.02	±	0.01	35.8	±	0.2
Maize	6.53	±	0.02	32.3	±	0.7
Banana	4.95	±	0.03	42.9	±	0.4
Pineapple	3.30	±	0.02	60.7	±	0.3
Yam	4.69	±	0.02	45.8	±	0.3
Tomato	4.16	±	0.02	51.2	±	0.8
Cassava	6.20	±	0.01	34.3	±	0.5
Grape	2.82	±	0.02	72.3	±	2.9
Mango	4.50	±	0.12	47.3	±	0.6
Orange	3.62	±	0.01	57.3	±	0.5

Note: Mean for three stations i.e., ST I: Akparabong in Ikom LGA, ST II: Boje in Boki LGA, ST III: Ugep in Yakurr LGA

Table 2: Dry Season Nov-April Mean pH and conductivity in Siemens ± Standard Deviation

Sample	Mean P ^H	±	STD	Mean Conductivity	±	STD
Cucumber	4.40	±	0.09	27.4	±	1.0
Plantain	4.62	±	0.12	24.2	±	4.1
Rice (white)	6.20	±	0.20	22.2	±	1.2
Maize	6.42	±	0.12	20.9	±	1.4
Banana	4.57	±	0.15	23.8	±	1.4
Pineapple	3.48	±	0.12	40.8	±	1.0
Yam	4.34	±	0.16	24.6	±	0.5
Tomato	4.25	±	0.06	25.6	±	0.6
Cassava	6.04	±	0.30	24.0	±	2.2
Grape	2.22	±	0.10	27.9	±	1.8
Mango	4.64	±	0.16	27.0	±	0.1
Orange	3.62	±	0.16	36.9	±	0.1
Cocoyam	4.85	±	0.21	25.3	±	3.5
Potato	4.38	±	0.08	26.5	±	0.2
Akpu (process cassava)	3.63	±	0.21	28.9	±	0.6
Cassava flour	9.37	±	0.14	33.8	±	2.9
Garri	4.64	±	0.16	23.7	±	0.2
Garri	4.13	±	0.13	25.7	±	0.2
Paw paw Sugar cane	3.42	±	0.12	32.3	±	1.0

Note: Mean for three stations i.e., ST I: Akparabong in Ikom LGA, ST II: Boje in Boki LGA, ST III: Ugep in Yakurr LGA

the consumption of carbohydrate source food items dislocates the pH balance of the body system and gives rise to innumerable health problems (Cleave, 1974; Susan, 2000; Young and Young, 2002). The rise in acidity is exacerbated by the hydrolysis of the food substances which are consumed by our local communities on a daily bases. This effect can be reduced either by the combination of other non-carbohydrate food substances in their appropriate quantities such as the proteins or by the increased in take of vegetables and fruits or by the fortification of these carbohydrate food substances with acids decreasing food substances or vitamins. A case in point is the treatment of raw cassava with a pH of 6.2 and Akpu (cassava meal) of pH of 3.6 to cassava flour with a pH of 9.37. Government and Non-Governmental agencies can be advised to carry out a sensitization campaign in the rural populace on the effect of consuming too much acidic foods and ways of minimizing it. Also, it is possible for government to legislate to the food manufacturing companies on the acid levels of their products and monitor them to comply. This will improve

the health condition of the people especially the rural communities.

REFERENCES

- Achinewhu, S.C., L.I. Barber and I.O. Ijeoma, 1998. Physicochemical Properties and Garification (gari yield) of selected cassava cultivars in River State, Nigeria. *Plant Food for Human Nutr.*, 52: 133-140.
- Akinrele, A., 1976. Improvement of Cassava processing. *Proc. 1st Int. Congress Lond.Tech.London.* 633-644.
- Anon, 1962. pH Values of Food Products. *Food Eng.*, 34: 98-99.
- Bridges, M.A. and M.R. Mattice, 1939. Over two thousand estimations of the pH of representative foods. *Am. J. Digestive Diseases*, 9: 440-449.
- Cleave, T.L., 1974. *The Saccharine Disease, The Master Disease of Our Time*, Keats Publishing, CT.
- Eka, O.U., 1986. Studies on fermentation of cassava, some enzymes and micro-organism in the fermentation liquour and the effect of fermentation on Nutrient states of cassava product. *W.A. J. Biol. and Appl. Chem.*, 31: 3-12.

- Eneobong, H., 2001. Eating Right. Nutrition Composition of Some Protein Foods. 142-146.
- Green, J. and C.R. Kleeman, 1991. The role of bone in the regulation of systematic acid base balance. *Contrib. Nephrol.*, 91: 61-76.
- Hall, R.H., 1976. Food for Naught, The decline in Nutrition Books, New York.
- Halprin, M.L. and M.B. Goldstein, 1999. Fluid, Electrolyte and acid-base physiology: A problem-Based Approach, 3rd Ed. Philadelphia. W.B Saunders company.
- Harris, J.F., 1975. Hydrolysis and Dehydration Reactions for Utilizing Plant Carbohydrates. *Appl. Polymer Symposium*, 131-144.
- Krist, H. and H.H. Rump, 1988. Laboratory Manual for Examination of Waste and Soil. London: Weinheim.
- Osuala, E.C., 1980. Introduction to research methodology (2nd) Benin City. Illupeju Press.
- Oyenuga, A., 1968. Nigerian foods and feeding stuffs. Ibadan, University press. pp: 9.
- Remer, T. and F. Manz, 1994. Estimation of the renal net acid excretion by adults consuming diets containing variable amount of protein. *Am. J. Clin. Nutr.*, 59: 1356-1361.
- Susan Brown, E., 2000. Acid-Alkaline Balance and its Effects on the Bone Health, *Int. J. Integrative Med.*, 2: 1-4.
- Young, R.O. and S.R. Young, 2002. P^H Miracle Problems of over acidity. pp: 29-39. Werner Books Inc., New York.