

Some Anti-Nutritional and Mineral Contents of Extra – Cotyledonous Deposit of Pride of Barbados (*Caesalpinia pulcherrima*)

T.P. Prohp¹, I.G. Ihimire², A.O. Madusha², H.O. Okpala³, J.O. Erebor³ and C.A. Oyinbo⁴

¹Department of Medical Biochemistry, ²Department of Biochemistry, ³Department of Medical Laboratory Sciences, ⁴Department of Anatomy, Ambrose Alli University, P.M.B. 14, Ekpoma, Edo State, Nigeria

Abstract: The edible extra-cotyledonous deposit of Pride of Barbados (*Caesalpinia pulcherrima*) was investigated for some minerals and anti-nutritional factors. Results obtained, showed that while phytohaemagglutinin was absent, vanadium was very poor in the extra-cotyledon of this legume. However, oxalate, trypsin inhibitor, phytate and polyphenol were found to be present at 18.09 ± 2.29 , 0.28 ± 0.00 , 96.40 ± 0.20 and 65.00 ± 0.02 mg/100g respectively. The sample was also found to contain the following essential minerals: iron (3.32 ± 0.09), zinc (3.59 ± 0.01), copper (1.36 ± 0.01), sodium (1.51 ± 0.04), potassium (6.22 ± 0.48), calcium (83.37 ± 0.24), Phosphorus (28.75 ± 0.37), magnesium (14.65 ± 0.24) and manganese (11.52 ± 0.13) mg/kg DM. Extra-cotyledonous deposit of Pride of Barbados is a good source of calcium and may serve as a cheap single source of a considerable number of essential minerals.

Key words: Anti-nutrients, minerals, extra-cotyledon, pride of barbados

Introduction

Pride of Barbados (*Caesalpinia pulcherrima*) as a legume belongs to the family leguminosae (Elegbede, 1998; Prohp *et al.*, 2004) which are known to contain biologically active principles. These anti-nutrient factors have adverse nutritional and physiological effects. According to Ihekoronye and Ngoddy (1985) some of these active principles interfere with digestive processes thereby preventing efficient utilization of the legume protein. Sometimes they are capable of precipitating deleterious effects in man and animals, with manifestable toxicity ranging from severe reduction in food intake and nutrient utilization to profound neurological effects culminating in death (Osagie, 1998). It has also been reported that tropical legumes contain a more complex array of these substances than any other crop species (Osagie, 1998). Minerals form an integral part of functionally important organic compounds such as iron (Fe) in haemoglobin or zinc (Zn) in insulin (Delvin, 1997). They are essential for the normal functioning of muscles, heart, nerves and in the maintenance of body fluid composition among others (White *et al.*, 1973). Mineral deficiencies have manifested in forms of different disease conditions as goitre, rickets and one form of metabolic dysfunction or the other.

Extra-cotyledonous deposit of Pride of Barbados is the translucent coat over the cotyledon embedded under the green testa of the seed (Prohp and Maduemezia, 2004). Children and adults eat this part of the plant from time to time. However, it is not a staple food in Nigeria (Prohp *et al.*, 2004). Earlier studies on some functional properties, nutrient potentialities, ash and moisture contents have shown that this edible part of Pride of Barbados (extra –

cotyledon) may be a potential food supplement (Prohp and Maduemezia, 2004; Prohp and Alaiya, 2003). Recent report has shown that of all the legumes analyzed in Nigeria, so far, Pride of Barbados presumably has the lowest cyanide content in the extra – cotyledon as well as in other parts of the plant (Prohp *et al.*, 2004).

In this work therefore, the aim is to determine mineral and some anti-nutritional contents of extra-cotyledonous deposit of Pride of Barbados with the view of further understanding its usefulness.

Materials and Methods

Pride of Barbados (*Caesalpinia pulcherrima*) plants grown around Faculty of Natural Sciences, Ambrose Alli University, Ekpoma, Edo State, Nigeria were the sources of experimental specimen. The mature pods were dissected to reveal the seeds in lateral arrangements. About 500g were air dried, milled and sieved through 1mm diameter sieve of a Hammer Miller (Glen Creston 14-5805) before analysis.

Analytical procedure: The mineral constituents of air-dried material were determined after wet digestion with a mixture of nitric, sulphuric and perchloric acid using Atomic absorption spectrophotometer. Sodium and potassium contents were determined with flame photometer. (AOAC, 1975).

Determination of antinutrients: Trypsin inhibitor was extracted with 1M NaOH and determined spectrophotometrically at 280nm.

Oxalate was determined by the method of Ukpabi and Ejidoh, 1989). Phytate was determined by the method of Maga (1982) while polyphenol content was determined

Table 1: Some anti-nutritional contents of extra-cotyledonous deposit of Pride of Barbados (*Caesalpinia pulcherrima*)

Anti-nutritional factor	Mg/100
Oxalate	18.09 ± 2.29
Phytohaemagglutinin	ABS
Trypsin Inhibitor	0.28 ± 0.00
Phytate	96.40 ± 0.20
Polyphenol	65.00 ± 0.02

ABS = absent. Values are mean + S. E. M. of three separate determinations

Table 2: Mineral contents of extra-cotyledonous deposit of Pride of Barbados (*Caesalpinia pulcherrima*)

Minerals	Mg/kg DM
*Iron (Fe)	3.32 ± 0.09
*Zinc (Zn)	3.59 ± 0.01
*Copper (Cu)	1.36 ± 0.01
Cadmium	2.68 ± 0.00
Lead (Pb)	2.71 ± 0.14
*Sodium (Na)	1.51 ± 0.04
*Potassium (K)	6.22 ± 0.48
*Calcium (Ca)	83.37 ± 0.24
Vanadium	0.15 ± 0.01
Cobalt (Co)	11.26 ± 0.02
Nickel (Ni)	11.19 ± 0.18
*Phosphorus (P)	28.75 ± 0.37
*Magnesium (Mg)	14.65 ± 0.24
*Manganese (Mn)	11.52 ± 0.13

*Some essential minerals determined.

Values are mean ± S.E.M. of three separate determinations.

spectrophotometrically at 500nm using vanillin reagent on catechin substrate.

Phytohaemagglutinin determination: The agglutinability of a 3% rabbit red blood cell was used as a quantitative assessment of phytohaemagglutinin content. Haemagglutinin was extracted from a 40% w/v mixture of finely ground sample (425ug in phosphate buffered saline (PBS), pH 7.4). The mixture was shaken with a mechanical shaker for 5 minutes before filtering through cheesecloth. For determination, one drop of red blood cell suspension and one drop of PBS, was shaken in a circular depression and agglutinability recorded as weak (+) moderate (++) , strong (+++), positive or negative (-) (Ihimire, 1997;Crichton and Walker, 1985).

Results

Table 1 and 2 are the results of some anti-nutritional and mineral contents of extra-cotyledonous deposit of Pride of Barbados (*Caesalpinia pulcherrima*) determined. Phytohaemagglutinin was absent (Table 1) in extra – cotyledon of this legume which also has very poor amount of vanadium, and high calcium content (Table 2).

Discussion

The nutritional importance of a given food depends on the nutrient and anti-nutritional constituents (Aletor *et al.*, 1994). The values of phytate and polyphenol contents determined (Table 1) in extra-cotyledonous deposit of Pride of Barbados, were lower than 234.00± 3.60mg/100g DM (for phytate) and 15.10 ± 0.80mg/g DM (for polyphenol) as reported for raw lima beans and lima beans boiled for 160 minutes respectively (Egbe, and Akinyele, 1990). The level of oxalate recorded was below reported values for fresh (raw) samples of some tropical leafy vegetables like Manihot essulenta, Talinum triangulare, Crassocephalum biafaiae and Celosia argentea (Aletor and Adeogun, 1995). Phytate, polyphenol and oxalate affect bioavailability of composite nutrients. They complex with bivalent ions like Ca²⁺, Mg²⁺, Fe²⁺ and Zn²⁺ making them unavailable especially in monogastric animals (Aletor and Omodara, 1994).

Extra-cotyledonous extract did not record any agglutination with 3% red blood cells or rabbit erythrocyte. Haemagglutinin, a toxic protein, found in soyabeans was absent in extra-cotyledonous deposit of Pride of Barbados. It is capable of imparting bitter or unacceptable taste when present in legumes alongside other anti-nutritional factors e.g. tannins, anthocyanins and phytates (Elegbede, 1998). The level of trypsin inhibitor obtained was low compared to that reported for lfe Bimpe, IT84E-124 and TVX716. The values reported were 2.11, 5.26, 3.16 and 2.11 mg/g respectively (Aletor and Aladetimi, 1989). However, the level of trypsin inhibitor detected in the sample was comparable to that reported for some vicia genotypes (Aletor and Adeogun, 1995). Trypsin inhibitor in high un-tolerable limit lowers the digestibility of legume proteins.

A total of fourteen minerals were determined in extra-cotyledonous deposit of Pride of Barbados. Calcium was predominantly present at 83.37 ± 0.24 mg/g (Table 2). However, the levels of the minerals recorded were lower than was reported in some leguminous browse plants like *Cassia alata*, *Cajanus cajan*, and *Cassia nodosa*. (Aletor and Omodara, 1994).

In conclusion, extra-cotyledon of Pride of Barbados is a good source of calcium and also contains an appreciable number of some essential minerals. It may, therefore, be explored for inclusion as an addition to the existing nutritional supplements.

References

- Aletor, V.A. A.V. Goodchild, E.L. Moneim and A.M. Abd , 1994. Nutritional and anti-nutritional characteristics of selected vicia genotypes. *Anim. Feed Sci. Tec.*, 47: 125-139.
- Aletor, V.A. and O.O. Aladetimi, 1989. Compositional evaluation of some cowpea varieties and some under-utilized edible legumes in Nigeria. *Nahrung*, 33: 999-1007.

Prohp et al.: The edible extra-cotyledonous deposit of Pride of Barbados

- Aletor, V.A. and O.A. Omodara, 1994. Studies on some leguminous browse plants with particular reference to their proximate, mineral and some endogenous anti-nutritional constituents. *Anim. Feed Sci. Tec.*, 46: 343-348.
- Aletor, V.A. and O.A. Adeogun, 1995. Nutrient and anti-nutrients components of some tropical leafy vegetables. *Food Chem. J.*, 53: 375-379.
- AOAC, 1975. Official Method of Analysis. 12th (ed) by W. Horowitz, Washington D. C. Crichton, P. B. and Walker, J. W. 1985. Methods for the detection of haemagglutinins in aeromonas. *J. Med. Microbiol.*, 49: 273-277.
- Delvin, T.M., 1997. Principles of nutrition II: micronutrients. In: *Textbook of Biochemistry with clinical correlations*. 4th (ed). John Wiley and Son Inc. New York, 124: 1139.
- Egbe, I.A. and I.O. Akinyele, 1990. Effect of cooking on the anti-nutritional factors of lima beans (*Phaseolus lunatus*). *Food Chem.*, 35: 81-87.
- Elegbede, J.A., 1998. Legumes. In: *Nutritional Quality of Plant Foods*. (Ed) Ambik Press Benin City Nig., 53-83.
- Ihekoronye, A. and P. Ngoddy, 1985. Proteins: food quality control. In: *Integrated Food Sciences and Technology*. 3rd (ed) Macmillian Publishers, London, 28: 155-193.
- Ihimire, I.G., 1997. Dehydrocyanation of lima bean, (*Phaseolus lunatus*). 1997; M.Sc. Thesis, Ambrose Alli University; Chemistry Dept., Ekpoma.
- Maga, J.A., 1982. Phytate. Its chemistry, occurrence, food interactions, nutritional significance and methods of analysis. *J. Agri. Food Chem.*, 30: 1-7.
- Osagie, A.U., 1998. Anti-nutritional factors. In: *Nutritional Quality of plant foods*. (ed) Ambolk press Benin City Nig., 21: 244.
- Prohp, T.P. and C. Maduemezia, 2004. Carbohydrate, ash and moisture contents of extra-cotyledonous deposits of pride of Barbados (*Caesalpinia pulcherrima*). *Nig. J. Agri. Sci. and Forestry*, 1: 195-204.
- Prohp, T.P., E.A. Mendie, A.O. Madusha, S.C. Uzoaru, A. Aigbiremolen and P.C. Onyebuagu, 2004. Cyanide contents of Pride of Barbados (*Caesalpinia pulcherrima*) grown in different parts of Nigeria. *J. Med. Lab. Sci.*, 13: 29-32.
- Prohp, T.P. and H.T. Alaiya, 2003. Some functional properties and anti-nutritional factors of extra-cotyledonous deposits of Pride of Barbados (*Caesalpinia pulcherrima*). *Proceedings (15th Animal conference of BSN held in AAU, Ekpoma, 40-45.*
- Ukpabi, V.J. and J.I. Ejidoh, 1989. Effect of deep oil frying on the oxalate content and the degree of itching of cocoyams. (*Xanthosoma* and *colocassia* spp). Technical paper presented at the 5th annual conference of the Agriculture society of Nigeria. Federal University of Technology Owerri, Nigeria, 3-6 (September).
- White, A., P. Handler and E.L. Smith, 1973. Body fluids and specialized tissues In: *principles of Biochemistry*. 6th (ed.) McGraw Hill kogakusha Ltd, Tokyo, 902: 1159.